

STUDIES IN THE EUPATORIEAE (ASTERACEAE). LIII.

A NEW GENUS, RAULINOREITZIA.

R. M. King and H. Robinson  
Smithsonian Institution, Washington, D.C. 20560.

Two of the genera, Acanthostyles and Raulinoreitzia, that we refer to the Disynaphia Complex have pyramidal panicle inflorescences. The slightly reddish colored and sometimes inclined or pendant series of heads are reminiscent of the habit of some grasses. Of these two genera, Raulinoreitzia is totally distinctive by its unlobed leaves, short-papillose stylar appendages and enlarged blunt apical cells on the pappus setae. The genus is distinguished from all others with such enlarged blunt apical cells on the pappus setae by the five flowered heads and by the lack of hairs on the base of the style.

Raulinoreitzia R.M.King & H.Robinson, genus novum Asteracearum (Eupatorieae). Plantae suffrutescentes erectae laxe ramosae. Caules teretes. Folia opposita distincae petiolata, laminis ellipticis vel basi cuneatis margine serrulatis. Inflorescentiae paniculatae; pedicelli glabri. Involucra squamae imbricatae ca. 15-20 valde inaequilongae ca. 4-seriatae glabrae ex parte rufescentes interiores anguste oblongae; receptacula parum convexa glabra. Flores 5 in capitulo; corollae anguste infundibulares 5-lobatae extus glanduliferae intus glabrae, cellulis angustis parietibus sinuositis, lobis aequilateraliter triangularibus vel longioribus extus ad apicem paucis papillosis; filamenta antherarum in parte superiore angusta, cellulis quadratis vel elongatis, parietibus annulate ornatis, thecis basi vix hastatis, cellulis exothecialibus subquadratis, appendicibus antherarum oblongis; styli inferne non incrassati glabri, appendicibus linearibus breviter papillosis; achaenia prismaticae 4-5 costata fere glabra, punctis saepe in seriebus transversis; carpopodia parva cylindrica superne dense breviter fimbriata, parietibus tenuibus; pappus setiformis uniseriatus, setis ca. 30 scabris ad apicem dilatatis, cellulis apicalibus inflatis obtusis.

Species typica: Baccharis crenulata Sprengel

The genus is named after P. Raulino Reitz recently appointed Director of the Jardim Botânico in Rio de Janeiro. Dr. Reitz is well known for his previous work at the Herbario "Barbosa Rodrigues" in Itajaí, Santa Catarina, Brasil. In addition to his other contributions, Dr. Reitz has served as

editor of *Sellowia* and of the important series of *Flora Ilustrada Catarinense*.

Our studies of the genus indicate that it contains the following two species.

Raulinoreitzia crenulata (Sprengel) R.M.King & H.Robinson, comb. nov. Baccharis crenulata Sprengel, Syst. Veg. 3: 466. 1826. Argentina, Bolivia, Brazil, Peru, Paraguay.

Raulinoreitzia tremula (Hook. & Arn.) R.M.King & H.Robinson, comb. nov. Eupatorium tremulum Hook. & Arn., Hook. Comp. Bot. Mag. 1: 241. 1836. Argentina, Brazil, Uruguay.

#### Acknowledgement

This study was supported in part by the National Science Foundation Grant GB 20502 to the senior author.